



Clavaseptin

Clavaseptin is a palatable antibiotic tablet used in dogs and cats, an antibiotic containing a combination of amoxicillin and a beta-lactomase inhibitor (which overcomes bacteria which are penicillin resistant). In dogs it is mainly used for dental infections, and in cats for skin infections.

Active ingredient: Amoxicillin 40 mg Clavulanic acid 10 mg.

Use: the recommended dose of the product is 10 mg amoxicillin /2.5 mg clavulanic acid per kg body weight twice a day by the oral route in dogs and cats, i.e. 1 tablet per 4 kg body weight every 12 h, according to the table in the data sheet.

Side effects: Do not use in case of hypersensitivity to penicillins or other substances of the β -lactam group. Do not administer to gerbils, guinea pigs, hamsters, rabbits and chinchillas. Special precautions for use in animals. In animals with impaired liver and kidney function, the use of the product should be subject to a risk/benefit evaluation by the veterinary surgeon and the posology evaluated carefully. Caution is advised in the use of Clavaseptin in any other small pet (non food-producing) herbivores. Use of the product should be based on susceptibility testing. Inappropriate use of the product may increase the prevalence of bacteria resistant to amoxicillin/clavulanic acid. Use of the product should take into account official and local antimicrobial policies. Adverse reactions Vomiting and diarrhoea may be observed. Treatment may be continued depending on the severity of the undesirable effect observed and a benefit/risk evaluation by the veterinary surgeon. Hypersensitivity reactions (allergic skin reactions, anaphylaxis) may be observed. In these cases, administration should be discontinued and a symptomatic treatment given. Use during pregnancy and lactation. The safety of the product has not been established during pregnancy and lactation. Laboratory studies in rats have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. Use the product only accordingly to the benefit/risk assessment by the responsible veterinarian. Interaction with other medicinal products and other forms of interaction. The bactericidal activity of amoxicillin may be reduced by the simultaneous use of bacteriostatic substances such as macrolides, tetracyclines, sulfonamides and chloramphenicol. Overdose (symptoms, emergency procedures, antidotes), if necessary.

Izplatītājs: SIA "Vet Line", Mazā Rāmavas iela 2, Valdlauči, +371 676 008 85 / +371 2026 9055